

JOSLIN DIABETES CENTER & JOSLIN CLINIC
CLINICAL NUTRITION GUIDELINE FOR OVERWEIGHT AND OBESE ADULTS WITH TYPE 2
DIABETES, PREDIABETES OR THOSE AT HIGH RISK FOR DEVELOPING TYPE 2 DIABETES
08 07 2011

The Joslin Clinical Nutrition Guideline For Overweight and Obese Adults With Type 2 Diabetes, Prediabetes or at High Risk for Developing Type 2 Diabetes is designed to assist primary care physicians, specialists, and other healthcare providers in individualizing the care of and set goals for adult, non-pregnant patients with type 2 diabetes or individuals at high risk for developing type 2 diabetes. This guideline focuses on the unique needs of those individuals, and complements the 2010 Dietary Guidelines for Americans, which is jointly developed by the Department of Health and Human Services and the Department of Agriculture. It is not intended to replace sound medical judgment or clinical decision-making and may need to be adapted for certain patient care situations where more or less stringent interventions are necessary.

The objectives of the Joslin Clinical Diabetes Guidelines are to support clinical practice and to influence clinical behaviors in order to improve clinical outcomes and assure that patient expectations are reasonable and informed. Guidelines are developed and approved through the Clinical Oversight Committee that reports to the Joslin Clinic Medical Director of Joslin Diabetes Center. The Clinical Guidelines are established after careful review of current evidence, medical literature and sound clinical practice. These Guidelines will be reviewed periodically and the Joslin Diabetes Center will maintain, upgrade or downgrade the rating for each recommendation when new evidence mandates such changes.

Joslin's Guidelines are evidence-based; in order to allow the user to evaluate the quality of the evidence used to support each standard of care, a modification of the GRADE system has been adopted. The table provided on page 5 describes the categories in which methodological quality and strength of recommendations have been classified. Evidence levels are graded 1A through 2C, as indicated in brackets.

Target Individuals and General Goals of Clinical Nutrition Guideline

*Target Population	BMI > 25 kg/m ² or Waistline > 40"/102 cm (men) [1B] > 35"/88 cm (women)	and or	Type 2 Diabetes Prediabetes IGT (impaired glucose tolerance) [1A] IFG (impaired fasting glucose) or High Risk for The Metabolic Syndrome (AHA/NHLBI criteria) [1B] Type 2 Diabetes Family history of type 2 DM (first degree relative) Confirmed diagnosis of insulin resistance (e.g., high basal insulin)
* For Asian populations (South Asian Indians, East Asians and Malays) a BMI >23 kg/m ² and a waistline >35"/90 cm in men or >31"/80 cm in women is considered. [1B]			

General Guidelines

- There is strong evidence that weight reduction improves insulin sensitivity and glycemic control, lipid profile, and blood pressure in type 2 diabetes and decreases the risk of developing type 2 diabetes in pre-diabetes and high-risk populations.
- To select an approach for medical nutrition therapy (MNT), target individuals should be referred to a registered dietitian (RD) or a qualified healthcare provider for assessment and review of medical management and treatment goals. [1B]
- Priorities of MNT for this population include:
 1. Weight reduction
 2. Meal to meal consistency in carbohydrate distribution for those with fixed medication/insulin programs
 3. Consideration of other nutrition related co-morbidities such as hypertension and dyslipidemia
- The meal plan composition, described below, is for general guidance only and may be individualized by the RD or other healthcare provider according to clinical judgment, individual (patient) preferences and needs, and metabolic response. The plan should be re-evaluated and modified to respond to changes in parameters such as blood pressure, A1C and frequency of hypo/hyperglycemia. Modification of goals may be needed for those requiring additional dietary considerations such as those with hyperkalemia or who are vegetarian.

Weight Reduction

1. A structured lifestyle plan that combines dietary modification, activity, and behavioral modification is necessary for weight reduction. [1B]
 2. A modest and gradual weight reduction of one to two pounds every one to two weeks should be the optimal target. [2A]. Reduction of daily caloric intake should be by range between 250 - 500 calories. [1C] Total daily caloric intake should not be less than 1000-1200 for women and 1200-1600 for men, or based on a RD assessment of usual intake. [1C]
 3. A 5-10% weight loss may result in significant improvement in blood glucose control among patients with diabetes and help prevent the onset of diabetes among individual with pre-diabetes. Weight reduction should be individualized and continued until an agreed upon BMI and/or other metabolic goals are reached. [2B]
 4. Target individuals should meet with RD to learn and practice portion control as an effective way of weight management. [1B]
 5. Meal replacements (MR)** in the form of shakes, bars, ready-to-mix powders, and pre-packaged meals that match these nutrition guidelines may be effective in initiating and maintaining weight loss [2 B]
 - Meal replacements should be used under the supervision of a RD.
 - When meal replacements are initiated, glucose levels should be carefully monitored and if needed, antihyperglycemic medications should be adjusted
- ** meal replacements should be used with caution by those with hyperkalemia
7. Bariatric surgeries, although not without medical and nutrition risks, are effective options and may be discussed when indicated (consider in individuals with BMI >40 kg/m² and those with BMI >35 kg/m² with other comorbidities). [2B]. To date, there is limited evidence to support the recommendation of bariatric surgeries for patients with BMI <35 kg/m² even if they have diabetes or other co-morbid conditions.

Macronutrient Composition

Fat	Percentage	There is general agreement that fat quality rather than quantity is important. The total fat intake should be generally limited to less than 35 % of total daily caloric intake [2B] <ul style="list-style-type: none"> • Saturated fat should be limited to < 7% of total caloric intake.[1B] • Polyunsaturated and monounsaturated fats should comprise the rest of the fat intake [2B] • Cholesterol limited to <300 mg/day or <200 mg/day in individuals with LDL-Cholesterol >100 mg/dl. [1C]
	Recommended	Mono and polyunsaturated fats (e.g., olive oil, canola oil, nuts/seeds, avocado and fish, particularly those high in omega-3 fatty acids). 4 oz of oily fish (e.g., salmon, herring, trout, sardines, fresh tuna) 2 times/week, as a source of omega-3 fatty acids. [1B]
	Not Recommended	Foods high in saturated fat, including beef, pork, lamb and high-fat dairy products (e.g., cream cheese, whole milk or yogurt) Foods high in trans-fats (e.g., fast foods, commercially baked goods, some margarines) Foods high in dietary cholesterol such as egg yolks, and organ meats.
Protein	Grams/day	Protein intake should not be less than 1.2 gm/kg of adjusted body weight Adjusted Body Weight = IBW (Ideal Body Weight) + 0.25 (Current Weight - IBW). This amount generally accounts for 20-30% of total caloric intake [2B] There are no reliable scientific findings to support a protein intake that exceeds 2 gm/kg of adjusted body weight. Emerging data suggest that protein aids in the sensation of fullness (low-protein meal plans are associated with increased hunger). A modest increase in protein reduces appetite and assists in achieving and maintaining weight reduction. [2B] Protein also helps to minimize loss of lean body mass. [2 B]
	Recommended	Fish, skinless poultry, nonfat or low-fat dairy, nuts, seeds, and legumes [2B]
	Not Recommended	High saturated fat protein sources in excess (e.g., beef, pork, lamb and high-fat dairy products), as they may be associated with increased cardiovascular risk.
	Patients with Renal Issues	Although reducing total calories may result in a reduction of the absolute total amount of protein intake, patients with signs of kidney disease (i.e., one or more of the following: proteinuria, GFR<60 ml/min) should consult a nephrologist before increasing total or percentage protein in their diet. [1B] Protein intake for these patients should be modified, but not lowered to a level that may jeopardize their overall health or increase their risk for malnutrition or hypoalbuminemia.

Macronutrient Composition (continued)

Carbohydrate	Percentage	<ul style="list-style-type: none"> Intake should be adjusted to meet the cultural and food preferences of the individual. The total daily intake of carbohydrate should be at least 130 gm/day and ideally 40-45% of the total caloric intake[1C]
	Consideration of Glycemic Index/Glycemic Load	The glycemic index/glycemic load is an important factor that patients should apply in their daily selection of carbohydrates foods. Foods with a low glycemic index should be selected [2B] (e.g., whole grains, legumes, fruits, green salad with olive oil-based dressing, and most vegetables)
	Recommended	Vegetables and fruits, legumes, whole and minimally processed grains. [2B]
	Not Recommended	Sugar, refined carbohydrates or processed grains and starchy foods especially sugary beverages, most pastas, white bread, white rice, low-fiber cereal and white potatoes should be consumed in limited quantities. [2B]
	Fiber	<ul style="list-style-type: none"> Approximately 14gm of fiber /1000 cal (20-35 gm) per day is recommended. [1B] If tolerated, ~50 gm/day is effective in improving postprandial hyperglycemia and should be encouraged. [2A] Fiber from unprocessed food, such as vegetables, fruits, seeds, nuts, and legumes is preferable but, if needed, fiber supplements such as psyllium, resistant starch and β-glucan can be added. [1B]

Micronutrient Composition

	Sodium	<ul style="list-style-type: none"> Daily consumption should be < 2300 mg (about 1 tsp of salt) per day. (1A) Further reduction to 1500 mg is recommended in people > 50 yr of age including those with chronic kidney disease. [2B] Slow acclimatization to lower sodium intakes is advisable.
	Potassium	<ul style="list-style-type: none"> Daily consumption should be a minimum of 4,700 mg unless potassium excretion is impaired Potassium helps offset high sodium intake by triggering more sodium excretion by the kidneys. Potassium-rich foods include fruits and vegetables like bananas, mushrooms, spinach, and almonds.

Dietary Supplements

In individuals who are not deficient, data do not support the use of vitamins or minerals to improve glucose control or the use of herbal supplements or spices to improve glucose control.

Non-nutritive Sweeteners

All FDA- approved non-nutritive sweeteners are permissible in moderate quantities (e.g., one diet soda daily)

Alcohol

- If consumed, alcohol consumption must be moderate. No more than 1 drink a day for women and no more than 2 drinks a day for men (one drink is equal to 12 ounces of regular beer, 5 ounces of wine, or 1.5 ounces of 80-proof distilled alcohol). [2C]
- Alcoholic beverages contain calories and are low in nutritional value.
- It is not advisable to increase alcohol consumption for the purpose of deriving purported health benefit.

Physical Activity and Behavioral Modification

- Physical activity should be included in the nutrition prescription described above. Increased physical activity should be an integral component of any weight reduction plan as it helps maximize the benefits of weight reduction on diabetes control and may prevent coronary and cerebral vascular disease. [1B]
- 60-90 minutes of moderately intensive activity, at least 5 days of the week, is encouraged for weight loss, unless contraindicated. [1B]
- Physical activity should be a mix of cardiovascular, flexibility, and resistance training to maintain or increase lean body mass.

Appendix A

Suggested Approximate Macronutrient Distribution According to Clinical Guideline

Calorie Level	Carbohydrate		Protein		Fat	
	Grams	%	Grams	%	Grams	%
1000	130	~50 *	75	30	27	20
1200	135	45	75-90	25-30	40	30
1500	150-170	40-45	75-110	20-30	50	30
1800	180-200	40-45	90-135	20-30	60	30
2000	200-225	40-45	100-150	20-30	70	~30

*A minimum of 130grams of carbohydrate per day, in a 1000 calorie meal plan, calculates to ~50% of the total daily calories.

Approved by the Joslin Clinical Oversight Committee on 08 07 2011

Clinical Nutrition Task Force	
Om Ganda, MD Co-Chair	Richard Jackson, MD
Osama Hamdy, MD, PhD-Co-Chair	Sharon Jackson, MS, RD, CDE
Gillian Arathuzik, MS, RD, CDE	Amanda Kirpitch, RD, CDE
Elizabeth Blair, MSN, ANP-BC, CDE	Melinda Maryniuk, MEd, RD, CDE
Cathy Carver, MSN, ANP-BC, CDE	Jo-Anne Rizzotto, MEd, RD, CDE
Amy Campbell, MS, RD, CDE	Nuha El Sayed, MD
Aaron Cypess, MD, PhD	Nora Saul, MS, RD, CDE
Edward S. Horton, MD	

Joslin Clinical Oversight Committee	
Om Ganda, MD -Chairperson	Melinda Maryniuk, MEd, RD, CDE
Richard Beaser, MD	Jo-Anne Rizzotto, Med, RD, CDE
Elizabeth Blair, MSN, ANP-BC, CDE	Bijan Roshan, MD
Amy Campbell, MS, RD, CDE	Susan Sjostrom, JD
Cathy Carver, MSN, ANP-BC, CDE	Kenneth Snow, MD
Jerry Cavallerano, OD, PhD	William Sullivan, MD
William Hsu, MD	Howard Wolpert, MD
Richard Jackson, MD	John Zrebiec, LICSW, CDE
Lori Laffel, MD, MPH	Martin Abrahamson, MD (<i>ex officio</i>)
Medha Munshi, MD	

Grading System Used in Guideline

Grade of Recommendation	Clarity of risk/benefit	Quality of supporting evidence
1A Strong recommendation High quality of evidence	Benefits clearly outweigh risk and vice versa.	Consistent evidence from well performed randomized, controlled trials or overwhelming evidence of some other form. Further research is unlikely to change our confidence in the estimate of benefit and risk.
1B Strong recommendation Moderate quality of evidence	Benefits clearly outweigh risk and burdens, or vice versa.	Evidence from randomized, controlled trials with important limitations (inconsistent results, methodological flaws, indirect or imprecise), or very strong evidence of some other research design. Further research is likely to have an impact on our confidence in the estimate of the benefit and risk and may change the estimate.
1C Strong recommendation Low quality of evidence	Benefits outweigh risk and burdens, or vice versa.	Evidence from observational studies, unsystematic clinical experience, or from randomized controlled trials with serious flaws. Any estimate of effect is uncertain.
2A Weak recommendation High quality of evidence	Benefits closely balanced with risks and burdens.	Consistent evidence from well performed randomized controlled trials or overwhelming evidence of some other form. Further research is unlikely to change our confidence in the estimate of benefit and risk.
2B Weak recommendation Moderate quality of evidence	Benefits closely balanced with risks and burdens; some uncertainty in the estimates of benefits, risks and burdens.	Evidence from randomized controlled trials with important limitations (inconsistent results, methodological flaws, indirect or imprecise), or very strong evidence of some other research design. Further research is likely to have an impact on our confidence in the estimate of benefit and risk and may change the estimate.
2C Weak recommendation Low quality of evidence	Uncertainty in the estimates of benefits, risks and burdens; benefits may be closely balanced with risks and burdens.	Evidence from observational studies, unsystematic clinical experience, or from randomized controlled trials with serious flaws. Any estimate of effect is uncertain.

Evidence graded less than “A” is acceptable to support clinical recommendations in a guideline. It is also assumed that for many important clinical recommendations, it would be unlikely that level A evidence be obtained because appropriate studies may never be performed.

¹ Guyatt G et al. Grading strength of recommendations and quality of evidence in clinical guidelines: Report from an American College of Physicians Task Force. *Chest* 129:174-181, 2006.

References.

1. Anderson JW, Gustafson NJ, Bryant CA, Tietyan-Clark J. Dietary fiber and diabetes: A comprehensive review and practical application. *J Am Diet Assoc*, 1987; 87(9):1189-97.
2. Andrews RC, Cooper AR, Montgomery AA, et al. Diet or diet plus physical activity versus usual care in patients with newly diagnosed type 2 diabetes: the Early ACTID randomised controlled trial. *Lancet* 2011; published online June 25. DOI:10.1016/S0140-6736(11)60442-X.
3. Baba NH, Sawaya S, Torbay N, Habbal Z, Azar S, Hashim SA. High protein vs high carbohydrate hypoenergetic diet for the treatment of obese hyperinsulinemic subjects. *Int J Obes Relat Metab Disord*, 1999;23(11):1202-1206.
4. Biesalski HK. Diabetes preventive components in the Mediterranean diet. *Eur J Nutr*, 2004;43 Suppl 1:I/26-30.
5. Brehm BJ, Seeley RJ, Daniels SR, D'Alessio DA. A randomized trial comparing a very low carbohydrate diet and a calorie-restricted low fat diet on body weight and cardiovascular risk factors in healthy women. *J Clin Endocrinol Metab*, 2003;88:1617-23.
6. Brinkworth GD, Noakes M, Parker B, Foster P, Clifton PM. Long-term effects of advice to consume a high-protein, low-fat diet, rather than a conventional weight-loss diet, in obese adults with type 2 diabetes: one-year follow-up of a randomised trial. *Diabetologia*, 2004;47(10):1677-86.
7. Chandalia M, Garg A, Lujthmann D, von Bergmann K, Grundy SM, Brinkley LJ. Beneficial effects of high dietary fiber intake in patients with type 2 diabetes mellitus. *N Engl J Med*, 2000;342:1392-1398.
8. Dansinger ML, Gleason JA, Griffith JL, Selker HP, Schaefer EJ. Comparison of the Atkins, Ornish, Weight Watchers, and Zone diets for weight loss and heart disease risk reduction: a randomized trial. *JAMA*, 2005;293(1):43-53.
9. DeCaterina Drug therapy: n-3 fatty acids in cardiovascular disease. *NEJM* 2011; 364: 2439-2450
10. Diabetes Prevention Program Research Group. Reduction in the Incidence of Type 2 Diabetes with Lifestyle Intervention or Metformin. *N Eng J Med*, 2002;346(6):393-403.
11. Dixon JB, O'Brien PE. Health outcomes of severely obese type 2 diabetic subjects 1 year after laparoscopic adjustable gastric banding. *Diabetes Care*, 2002;25(2):358-63.
12. Dumesnil JG, Turgeon J, Tremblay A, Poirier P, Gilbert M, Gagnon L, St-Pierre S, Garneau C, Lemieux I, Pascot A, Bergeron J, Despres JP. Effect of a low-glycaemic index--low-fat--high protein diet on the atherogenic metabolic risk profile of abdominally obese men. *Br J Nutr*, 2001;86(5):557-68.
13. Ferchak CV, Meneghini LF. Obesity, bariatric surgery and type 2 diabetes--a systematic review. *Diabetes Metab Res Rev*, 2004;20(6):438-45.
14. Foster GD, Wyatt HR, Hill JO, Makris AP, Rosenbaum DL, Brill C, Stein RI, Mohammed BS, Miller B, Rader DJ, Zemel B, Wadden TA, Tenhave T, Newcomb CW, Klein S. Weight and metabolic outcomes after 2 years on a low-carbohydrate versus low-fat diet: a randomized trial. *Ann Intern Med*. 2010;153(3):147-57
15. Foster GD, Wyatt HR, Hill JO, McGuckin BG, Brill C, Mohammed BS, Szapary PO, Rader DJ, Edman JS, Klein S. A randomized trial of a low-carbohydrate diet for obesity. *N Engl J Med*, 2003;348(21):2082-90.
16. Fukagawa NK, Anderson JW, Hageman G, Young VR, Minaker KL. High-carbohydrate, high-fiber diets increase peripheral insulin sensitivity in healthy young and old adults. *Am J Clin Nutr*, 1990;52:524-8.
17. Gannon MC, Nuttall FQ, Saeed A, Jordan K, Hoover H. An increase in dietary protein improves the blood glucose response in persons with type 2 diabetes. *Am J Clin Nutr*, 2003;78(4):734-41.
18. Gannon MC, Nuttall FQ. Effect of a high-protein, low-carbohydrate diet on blood glucose control in people with type 2 diabetes. *Diabetes*, 2004;53(9):2375-82.
19. Hamdy O, Horton ES. Protein Content in Diabetes Nutrition Plan. *Curr Diab Rep*. 2011 Jan 5. [Epub ahead of print]
20. Hamdy O, Ledbury S, Mullooly C, Jarema C, Porter S, Ovalle K, Moussa A, Caselli A, Caballero AE, Economides PA, Veves A, Horton ES. Lifestyle modification improves endothelial function in obese subjects with the insulin resistance syndrome. *Diabetes Care*, 2003; 26(7): 2119-25.
21. Hamdy, O, Morsi,A, El-Sayed,N et al Long-term weight reduction in clinical practice after non- surgical diabetes weight management program. *Diabetes* 2011, 60(suppl1): A515
22. Hu FB, Stamper MJ, Manson JE, et al. Dietary fat intake and the risk of coronary heart disease in women. *N Engl J Med*, 1997;337:1491-9.
23. Hu FB. Plant-based foods and prevention of cardiovascular disease: an overview. *Am J Clin Nutr*, 2003;78(3 Suppl):544S-551S.
24. Johnston CS, Tjonn SL, Swan PD. High-protein, low-fat diets are effective for weight loss and favorably alter biomarkers in healthy adults. *J Nutr*, 2004;134(3):586-91.
25. Keno Y, Matsuzawa Y, Tokunaga K, Fujioka S, Kawamoto T, Kobatake T, Tarui S. High sucrose diet increases visceral fat accumulation in VMH-lesioned obese rats. *Int J Obes*, 1991;15(3):205-11.
26. Kim JY, Nolte LA, Hansen PA, Han DH, Ferguson K, Thompson PA, Holloszy JO. High-fat diet-induced muscle insulin resistance: relationship to visceral fat mass. *Am J Physiol Regul Integr Comp Physiol*, 2000;279(6):R2057-65.
27. Lara-Castro, C, Garvey, T. Diet, insulin resistance, and obesity: Zoning in on data for Atkins dieters living in South Beach. *J Clin Endocrinol Metab*, 2004;89:4197-4205.
28. Larsen TM, Dalskov SM, van Baak M, Jebb SA, Papadaki A, Pfeiffer AF, Martinez JA, Handjieva-Darlenska T, Kunešová M, Pihlsgård M, Stender S, Holst C, Saris WH, Astrup A; Diet, Obesity, and Genes (Diogenes) Project. Diets with high or low protein content and glycemic index for weight-loss maintenance. *N Engl J Med*. 2010;363(22):2102-13.
29. Look AHEAD Research Group. Long-term effects of a lifestyle intervention on weight and cardiovascular risk factors in individuals with type 2 diabetes mellitus: four-year results of the Look AHEAD trial. *Arch Intern Med* 2010; 170: 1566-75.
30. Ludwig DS, Pereira MA, Kroenke CH, et al. Dietary fiber, weight gain, and cardiovascular disease risk factors in young adults. *JAMA*, 1999; 282:1539-46.
31. Ludwig DS. Dietary glycemic index and obesity. *J Nutr*, 2000;130:280S-3S.
32. Luscombe ND, Clifton PM, Noakes M, Parker B, Wittert G. Effects of energy-restricted diets containing increased protein on weight loss, resting energy expenditure, and the thermic effect of feeding in type 2 diabetes. *Diabetes Care*, 2002;25(4):652-7.
33. Mann JI. Diet and risk of coronary heart disease and type 2 diabetes. *Lancet*, 2002;360(9335):783-9.
34. Meckling KA, O'Sullivan C, Saari D. Comparison of a low-fat diet to a low-carbohydrate diet on weight loss, body composition, and risk factors for diabetes and cardiovascular disease in free-living, overweight men and women. *J Clin Endocrinol Metab*, 2004;89(6):2717-23.

35. Misra A, Wasir JS, Pandey RM. An evaluation of candidate definitions of the metabolic syndrome in adult Asian Indians. *Diabetes Care* 2005;28(2):398-403.
36. McAuley KA, Hopkins CM, Smith KJ, McLay RT, Williams SM, Taylor RW, Mann JI. Comparison of high-fat and high-protein diets with a high-carbohydrate diet in insulin-resistant obese women. *Diabetologia*, 2005;48(1):8-16.
37. Park, Y, Subar, AF, Hollenbeck, A Schatzkin, A Dietary fiber intake and mortality in the NIH-AARP Diet and Health study. *Arch Intern Med* 2011; 171:1061-1068
38. Parker B, Noakes M, Luscombe N, Clifton P. Effect of a high-protein, high-monounsaturated fat weight loss diet on glycemic control and lipid levels in type 2 diabetes. *Diabetes Care*, 2002;25(3):425-30.
39. Pereira MA, Swain J, Goldfine AB, Rifai N, Ludwig DS. Effects of a low glycemic load diet on resting energy expenditure and heart disease risk factors during weight loss. *JAMA* 2004;292:2482-90.
40. Rock CL, Flatt SW, Sherwood NE, Karanja N, Pakiz B, Thomson CA. Effect of a free prepared meal and incentivized weight loss program on weight loss and weight loss maintenance in obese and overweight women: a randomized controlled trial. *JAMA*. 2010;304(16):doi:10.1001/jama.2010.1503
41. Skov AR, Toubro S, Ronn B, Holm L, Astrup A. Randomized trial on protein vs carbohydrate in ad libitum fat reduced diet for the treatment of obesity. *Int J Obes Relat Metab Disord*, 1999;23(5):528-36.
42. Sofi, F Cesari, F, Abbate, R et al Adherence to Mediterranean diet and health status: a meta analysis. *BMJ* 2008; 337: a1344
43. Stern L, Iqbal N, Seshadri P, Chicano KL, Daily DA, McGrory J, Williams M, Gracely EJ, Samaha FF. The effects of low-carbohydrate versus conventional weight loss diets in severely obese adults: one-year follow-up of a randomized trial. *Ann Intern Med*, 2004;140(10):778-85.
44. Story L, Anderson JW, Chen WJ, Karounos D, Jefferson B. Adherence to high-carbohydrate, high-fiber diets: Long-term studies of non-obese diabetic men. *J Am Diet Assoc*, 1985;85(9):1105-10.
45. Suzukawa M, Abbey M, Howe PR, Nestel PJ. Effects of fish oil fatty acids on low density lipoprotein size, oxidizability, and uptake by macrophages. *J Lipid Res*, 1995;36(3):473-84.
46. Uusitupa M, Lindi V, Louheranta A, Salopuro T, Lindstrom J, Tuomilehto J; Finnish Diabetes Prevention Study Group. Long-term improvement in insulin sensitivity by changing lifestyles of people with impaired glucose tolerance: 4-year results from the Finnish Diabetes Prevention Study. *Diabetes*, 2003;52(10):2532-8.
47. Yang, Q Liu, T, Kuklina, EV et al Sodium and potassium intake and mortality among US adults. Prospective data from Third NHANES. *Arch Intern Med* 2011; 171: 1183-1191
48. Yip I, Go VL, DeShields S, Saltsman P, Bellman M, Thames G, Murray S, Wang HJ, Elashoff R, Heber D. Liquid meal replacements and glycemic control in obese type 2 diabetes patients. *Obes Res*, 2001;9 Suppl 4:341S-347S..